

- I. (a) (i) I. Complete the expression given below to determine the charge ( $Q$ ) of an atom in a Lewis structure by inserting the terms  $N_A$ ,  $N_{LP}$  and  $N_{BP}$  in the appropriate boxes, where.

$N_A$  = number of valence electrons in the atom

$N_{LP}$  = number of electrons in lone pairs

$N_{BP}$  = number of electrons in bonding pairs around the atom

$$Q = \boxed{N_A} - \boxed{N_{LP}} - \frac{1}{2} \boxed{N_{BP}}$$

(01)                    (01)                    (01)

- II. Fill in the values for  $N_A$ ,  $N_{LP}$  and  $N_{BP}$  in the appropriate boxes and calculate the charge on S, Q(sulfur), in the structure  $SOP_2$  given below.




$$Q(\text{sulfur}) = \boxed{6} - \boxed{2} - \frac{1}{2} \boxed{8} = \dots \dots \dots$$

(01)                    (01)                    (01)                    (04)

Mark Independently



(10 marks)

- (ii) Draw the most acceptable Lewis structure for the ion,  $\text{ClO}_2\text{F}_2^-$ .



(07 marks)

- (iii) The most stable Lewis structure for the molecule  $\text{CH}_2\text{SO}$  (sulfine) is shown below. Draw another two Lewis structures (resonance structures) for this molecule.



Disregard the double headed arrows


(Any two)

(07 marks x 2 = 14 marks)

- (iv) Based on the hypothetical Lewis structure given below, state the following regarding the C, N and O atoms given in the table below.

- I. VSEPR pairs around the atom  
III. shape around the atom

- II. electron pair geometry around the atom  
IV. hybridization of the atom



The atoms are numbered as follows:



(v) Identify the atomic/hybrid orbitals involved in the formation of the following  $\sigma$  bonds in the Lewis structure given in part (iv) above. (Numbering of atoms is as in part (iv).)

I.	$\text{N}^2-\text{C}^3$	$\text{N}^2 \dots \text{sp}$	$\text{C}^3 \dots \text{sp}^2$
II.	$\text{O}^4-\text{N}^5$	$\text{O}^4 \dots \text{sp}^3$	$\text{N}^5 \dots \text{sp}^3$
III.	$\text{N}^5-\text{H}^6$	$\text{N}^5 \dots \text{sp}^3$	$\text{H}^6 \dots \text{1s}/\text{s}$
IV.	$\text{C}^7-\text{O}^7$	$\text{C}^7 \dots \text{sp}^2$	$\text{O}^7 \dots \text{sp}^2 \text{ OR } 2p$

(01 marks x 8 = 08 marks)

1(a): 55 marks

(b) (i) Identify the sub-shells (atomic orbitals) along with their azimuthal quantum number ( $l$ ), and magnetic quantum numbers ( $m_l$ ) for the energy level with principal quantum number  $n=3$  in an atom. What is the maximum number of electrons present in each sub-shell?

Write your answers in the table given below.

Sub-shell	Azimuthal quantum number ( $l$ )	Magnetic quantum numbers ( $m_l$ )	Maximum number of electrons in each sub-shell
$3s/s$	0	0	2
$3p/p$	1	-1, 0, +1	6
$3d/d$	2	-2, -1, 0, +1, +2	10

(01 marks x 12 = 12 marks)

(ii) Identify the type/s of intermolecular forces present in I, II and III given below.

I. Ar gas  
London dispersion forces

II. NO gas  
dipole-dipole + London dispersion forces

III. water sample containing a small amount of dissolved KCl  
Ion dipole + Hydrogen bonding

(01 marks x 5 = 05 marks)

(iii) "The boiling point of n-butane ( $\text{C}_4\text{H}_{10}$ ) is higher than the boiling point of propane ( $\text{C}_3\text{H}_8$ )."  
Giving reasons, state whether this statement is true or false.

True ..... (05)

n-butane and propane do not possess dipole moments or n-butane and propane are nonpolar molecules ..... (02)

Forces acting are London dispersion forces ..... (02)

Since n-butane is greater in size or molar mass than propane ..... (02)

London dispersion forces between n-butane molecules are greater than in propane molecules ..... (02)

Therefore, boiling point of  $\text{C}_4\text{H}_{10} > \text{C}_3\text{H}_8$  .....

(iv) Arrange the following in the decreasing order of the property indicated in parentheses. (Reasons are not required.)

I.  $\text{Li}_2\text{CO}_3$ ,  $\text{Na}_2\text{CO}_3$ ,  $\text{K}_2\text{CO}_3$  (solubility in water)

$\text{K}_2\text{CO}_3 > \text{Na}_2\text{CO}_3 > \text{Li}_2\text{CO}_3$  ..... (05)

II.  $\text{NF}_3$ ,  $\text{NH}_3$ ,  $\text{NOCl}$ ,  $\text{NO}_2^+$  (bond angle)

$\text{NO}_2^+ > \text{NOCl} > \text{NH}_3 > \text{NF}_3$  ..... (05)

III.  $\text{COCl}_2$ ,  $\text{CO}_2$ ,  $\text{HCN}$ ,  $\text{CH}_3\text{Cl}$  (electronegativity of carbon)

$\text{CO}_2 > \text{HCN} > \text{COCl}_2 > \text{CH}_3\text{Cl}$  ..... (05)

1(b): 45 marks

3. (a) X, Y and Z are elements that belong to the same group in the Periodic Table. They are in three successive periods respectively on descending the group. Y exists as a non-metallic coloured liquid at room temperature.

(i) Identify X, Y and Z. (Give atomic symbols.)

$$X = \dots Cl \dots \quad Y = \dots Br \dots \quad Z = \dots I \dots$$

Note: Do not award marks for X: Cl, Y: Br, Z: I.

(04 marks x 3 = 12 marks)

(ii) Indicate the relative magnitudes of the following with regard to X, Y and Z.

I. Atomic size

I	>	Br	>	Cl
Cl	>	Br	>	I
Cl	>	Br	>	I

II. Electron affinity

III. First ionization energy

Note: X, Y and Z have all to be identified correctly in order to award marks for 2(a)(ii) provided all three elements are identified correctly X, Y & Z can be used to fill boxes

(03 marks x 3 = 09 marks)

(iii) You are provided with aqueous solutions of the anions of X, Y and Z, each in a separate test-tube. Suggest a single reagent that could be used to identify these anions.

[Note: You are required to state the observation for each anion.]

Reagent:  $\text{AgNO}_3$  solution ..... (04)

Observation: (for the anions) X: white precipitate ..... (02)

Y: pale yellow precipitate ..... (02)

Z: dark yellow precipitate ..... (02)

OR

Reagent:  $\text{Cl}_2/\text{CCl}_4$  ..... (04)

Observation: (for the anions) X: colourless organic layer ..... (02)




Y: red-orange organic layer ..... (02)

Z: violet organic layer ..... (02)

Note: X, Y and Z have all to be identified correctly in order to award marks for 2(a)(iii)




Even if X, Y and Z have been identified as  $\text{Cl}_2$ ,  $\text{Br}_2$ , and  $\text{I}_2$  marks can be awarded

(iv) Give balanced chemical equations for the reactions of  $\text{Cl}_2(\text{g})$  with the following.



Note: Physical states are not required for the award of marks

(v) Draw the structures of two oxoacids of X.




Any two (Neglect shapes and lone pairs however all bonds must be shown)

) ..... (03 x 2 = 06 marks)

(vi) Name one natural source of X. Sea water/Rock salt ( $\text{NaCl}$ )/KCl (Sylvine)/KCl.MgCl<sub>2</sub> (Carnallite) ..... (02)

(vii) I. A monomer that contains X forms an addition polymer that is widely used in the manufacture of water pipes. Draw the structure of the monomer.



(03)

II. Write the full name of the polymer. Polyvinyl chloride ..... (02)

2(a): 50 marks


- (b) An aqueous solution Q contains three anions. The following tests were carried out to identify these anions.  
(Fresh portions of solution Q were used for each test ① to ⑤).

	Test	Observation
①	I Dilute HCl was added.	A colourless gas was evolved. A clear solution was obtained.
	II The gas evolved was tested with filter paper moistened with lead acetate.	No colour change
②	I A $\text{BaCl}_2$ solution was added.	A white precipitate was obtained.
	II The white precipitate was separated by filtration, and dil. HCl was added to it.	The white precipitate dissolved with the evolution of a gas.
	III The gas evolved was tested with a filter paper moistened with acidified potassium dichromate.	The colour changed from orange to green.
③	Conc. $\text{HNO}_3$ and an excess of ammonium molybdate solution were added and the mixture was warmed.	A yellow precipitate did not form.
④	Devarda's alloy and NaOH solution were added and the mixture was heated.	A gas that turned Nessler's reagent brown was evolved.
⑤	A $\text{FeCl}_3$ solution was added.	A blood red coloured solution was obtained.





- (i) Identify the three anions in solution Q.

$$\text{NO}_3^- \dots , \text{SCN}^- \dots \text{ and } \text{SO}_4^{2-} \dots \quad (12 \times 3 = 36 \text{ marks})$$

- (ii) Write the balanced chemical equation for the reaction taking place in test number ② III.




Or



2(b): 50 marks

3. (a) Methylamine,  $\text{CH}_3\text{NH}_2$  is a weak base. The following equilibrium exists in an aqueous solution of methylamine.




- (i) Write the expression for  $K_b$  of methylamine.

Note: Correct Physical states are necessary

$$K_b = \frac{[\text{CH}_3\text{NH}_3^+(\text{aq})][\text{OH}^-(\text{aq})]}{[\text{CH}_3\text{NH}_2(\text{aq})]} \quad (05)$$

- (ii) At 25°C, the pH value of a  $0.20 \text{ mol dm}^{-3}$  aqueous solution of methylamine is 11.00. Calculate  $K_b$ .



$$\text{As the pH} = 11, \text{ pOH} = 3 \quad (02)$$


$$\therefore [\text{OH}^-(\text{aq})] = [\text{CH}_3\text{NH}_3^+(\text{aq})] = 1.0 \times 10^{-3} \text{ mol dm}^{-3} \quad (02)$$

$$\therefore K_b = \frac{1.0 \times 10^{-3} \text{ mol dm}^{-3} \times 1.0 \times 10^{-3} \text{ mol dm}^{-3}}{0.2 \text{ mol dm}^{-3}} \quad (02 + 01)$$

$$\therefore = 5.0 \times 10^{-6} \text{ mol dm}^{-3} \quad (02 + 01)$$

- (iii) A volume of  $25.00 \text{ cm}^3$  of the solution in (ii) above was titrated with  $0.20 \text{ mol dm}^{-3}$  HCl at  $25^\circ\text{C}$ . Calculate the pH value of the solution at the equivalence point. ( $K_w = 1.0 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$  at  $25^\circ\text{C}$ .)

Reaction of  $\text{CH}_3\text{NH}_2(\text{aq})$  with HCl is,



Stoichiometry of the reaction,



Equivalence point =  $25.00 \text{ cm}^3$

(02)

$$\therefore [\text{CH}_3\text{NH}_3^+(\text{aq})] \text{ at the equivalence point} = \frac{0.2 \text{ mol dm}^{-3} \times 25 \times 10^{-3} \text{ dm}^3}{50 \times 10^{-3} \text{ dm}^3} = 0.10 \text{ mol dm}^{-3} \quad (04 + 01)$$

pH at the equivalence point is determined by the reaction;



<i>Initial con.</i>	0.10	0	0
<i>Egmt con.</i>	$0.10 - x$	x	$x \text{ mol dm}^{-3}$

(02)

For the Reaction (A):

$$K_a = \frac{[\text{CH}_3\text{NH}_2(\text{aq})][\text{H}_3\text{O}^+(\text{aq})]}{[\text{CH}_3\text{NH}_3^+(\text{aq})]} = \frac{K_w}{K_b} \quad (02)$$

$$= \frac{1.0 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}}{5.0 \times 10^{-6} \text{ mol dm}^{-3}} = 2.0 \times 10^{-9} \text{ mol dm}^{-3} \quad (03)$$

$$\therefore 2.0 \times 10^{-9} = \frac{x^2}{0.10 - x} \approx \frac{x^2}{0.10} \quad (02 + 03)$$

$$2.0 \times 10^{-10} = x^2$$


$$x = 1.41 \times 10^{-5} \text{ mol dm}^{-3} \quad (04 + 01)$$

$$\therefore \text{pH} = -\log(1.41 \times 10^{-5}) = 4.85 \quad (05)$$

3(a): 50 marks

- (b) In an experiment, a limited volume of  $1.00 \text{ mol dm}^{-3}$   $\text{HNO}_3$  was added to a precipitate  $\text{MX}(\text{s})$  and the system was allowed to reach equilibrium at  $25^\circ\text{C}$ . This resulted in partial dissolution of the precipitate giving rise to a clear solution. The  $\text{HX}(\text{aq})$  formed behaves as a weak acid.

- (i) Write chemical reactions for the equilibria existing in the above solution.



- (ii) Calculate  $[\text{X}^-(\text{aq})]$  in the solution assuming that the dissociation of  $\text{HX}(\text{aq})$  is negligible. (Solubility product of  $\text{MX}$  at  $25^\circ\text{C}$ ,  $K_{sp(\text{MX})} = 3.6 \times 10^{-7} \text{ mol}^2 \text{ dm}^{-6}$ .)



$$\text{The concentration of } \text{M}^+(\text{aq}) = 1.0 \text{ mol dm}^{-3} \quad (04 + 01)$$

$$K_{sp(\text{MX})} = [\text{M}^+(\text{aq})][\text{X}^-(\text{aq})] \quad (05)$$

$$\therefore [\text{X}^-(\text{aq})] = \frac{3.6 \times 10^{-7} \text{ mol}^2 \text{ dm}^{-6}}{1.0 \text{ mol dm}^{-3}} = 3.6 \times 10^{-7} \text{ mol dm}^{-3}$$

(04 + 01)

- (iii) Giving reasons explain whether  $[X^{-(aq)}]$  in a saturated aqueous solution of MX at 25°C is equal to, smaller than or greater than the value obtained in (ii) above.  
In an aqueous solution,  $[X^{-(aq)}]$  is determined only by  $K_{sp}$  (05)

$$K_{sp(MO)} = [M^{+}(aq)][X^{-(aq)}] = 3.6 \times 10^{-7} \text{ mol}^2 \text{ dm}^{-6}$$

$$[X^{-(aq)}]^2 = 3.6 \times 10^{-7} \text{ mol}^2 \text{ dm}^{-6} \quad (05)$$

$$[X^{-(aq)}] = 6.0 \times 10^{-4} \text{ mol dm}^{-3} \quad (05)$$

The value is higher than that in (ii) (05)

OR  $[X^{-(aq)}]$  is determined only by  $K_{sp}$  (05)

No common ion effect (05)




Solution is saturated (05)

The value is larger (05)

3(b): 50 marks

4. (a) The alcohols A, B, C and D are structural isomers of each other having the molecular formula  $C_5H_{12}O$ . A, B and C show optical isomerism.

- (i) Draw possible structures for A, B and C.




Structures can be drawn in any order

15-  $C_3H_7 \rightarrow$  only one mark.

When B, C and D are reacted with acidic  $K_2Cr_2O_7$ , X, Y and Z are formed respectively. The products X, Y and Z can be converted back to B, C and D respectively by reacting with  $\text{NaBH}_4$ .


- (ii) What is the structure of A?




A

On heating with conc.  $H_2SO_4$  A and B gave E and F, respectively, while C and D give the same product G. G shows diastereoisomerism. All three compounds E, F and G have the molecular formula  $C_5H_{10}$ . When E and F are reacted with HBr the same product H was formed.


- (iii) Draw the structures of B, C, D, E, F and H.




B




C




D



E




F



H

(iv) Draw the structures of the diastereoisomers of G.




Structures can be drawn in any order

04 marks X 12 = 48 marks


4(b): 48 marks

(b) Consider the two reaction schemes given below.




(i) Draw the structures of J, K and L in the boxes given below.


Note: If J =




03 marks only



J



K



L


Note: Award marks even if OH is written instead of O- in the structure K.

(ii) Write the reagents V and W in the boxes given below.


V = NaOH Or OH-W = PCC(iii) Writing A<sub>E</sub>, A<sub>N</sub>, S<sub>E</sub>, S<sub>N</sub> or E in the appropriate box, classify each of the reactions 1, 2 and 3 as electrophilic addition (A<sub>E</sub>), nucleophilic addition (A<sub>N</sub>), electrophilic substitution (S<sub>E</sub>), nucleophilic substitution (S<sub>N</sub>) or elimination (E) reaction.Reaction 1 S<sub>E</sub>Reaction 2 S<sub>N</sub>Reaction 3 A<sub>N</sub>

05 marks X 8 = 40 marks

4(b): 40 marks

(c) (i) What is the structure of the major product of the reaction between CH<sub>3</sub>CH=CH<sub>2</sub> and HBr?

(ii) Write the mechanism of the above reaction.



4(c): 12 marks

Note: Ignore δ+ and δ- if written near the H and Br of HBr and award marks.

5. (a) The following reaction occurs when  $\text{NaHCO}_3(s)$  is heated to a temperature above  $100^\circ\text{C}$ .



A sample of  $\text{NaHCO}_3(s)$  was placed in an evacuated closed rigid container of  $5.00 \text{ dm}^3$  volume and heated to  $328^\circ\text{C}$ . After the equilibrium was reached, a small amount of  $\text{NaHCO}_3(s)$  still remained in the container. The pressure of the container was found to be  $1.0 \times 10^6 \text{ Pa}$ . Assume that the volume of the solids remaining in the container is negligible.  $RT = 3000 \text{ J mol}^{-1}$  at  $328^\circ\text{C}$ .

(i) Calculate the number of moles of  $\text{H}_2\text{O}(g)$  in the container when the equilibrium is reached at  $328^\circ\text{C}$ .



System has  $\text{CO}_2(g)$  and  $\text{H}_2\text{O}(g)$  as gaseous species,

Assuming ideal behavior or ( $PV = nRT$ ) (05)

$$n_{\text{total}} = P_{\text{total}}V/RT$$

$$= 1.0 \times 10^6 \text{ Pa} \times 5.00 \times 10^{-3} \text{ m}^3 / 8.314 \text{ J K}^{-1} \text{ mol}^{-1} \times 601 \text{ K} \quad (02)$$

$$= 1.0 \times 10^6 \text{ Pa} \times 5.00 \times 10^{-3} \text{ m}^3 / 5000 \text{ J mol}^{-1}$$

$$= 1.0 \text{ mol} \quad (05)$$

$$n_{\text{total}} = n_{\text{H}_2\text{O}(g)} + n_{\text{CO}_2(g)}$$

According to stoichiometry  $n_{\text{H}_2\text{O}(g)} = n_{\text{CO}_2(g)}$  (02)

$$\therefore n_{\text{H}_2\text{O}(g)} = 0.50 \text{ mol} \quad (03)$$

(ii) Calculate  $K_p$  for the above equilibrium at  $328^\circ\text{C}$ , and hence calculate  $K_c$ .

$$K_p = P_{\text{H}_2\text{O}(g)} \times P_{\text{CO}_2(g)} \quad (05)$$

$$\text{in the system, } P_{\text{Total}} = P_{\text{H}_2\text{O}(g)} + P_{\text{CO}_2(g)} = 1.0 \times 10^6 \text{ Pa} \quad (05)$$

$$\text{and } P_{\text{H}_2\text{O}(g)} = P_{\text{CO}_2(g)}$$

$$\therefore P_{\text{H}_2\text{O}(g)} = P_{\text{CO}_2(g)} = 5.0 \times 10^5 \text{ Pa} \quad (04 + 01)$$

$$\therefore K_p = (5.0 \times 10^5 \text{ Pa})^2 = 2.5 \times 10^{11} \text{ Pa}^2 \quad (04 + 01)$$

$$K_p = K_c (RT)^{\Delta n} \quad (05)$$

$$\Delta n = 2 - 0 = 2 \quad (03)$$

$$\begin{aligned} \therefore K_c &= K_p / (RT)^2 \\ &= 2.5 \times 10^{11} \text{ Pa}^2 / (8.314 \text{ J K}^{-1} \text{ mol}^{-1} \times 601 \text{ K})^2 \\ &= 2.5 \times 10^{11} \text{ Pa}^2 / (5000 \text{ J mol}^{-1})^2 \\ &= 1.0 \times 10^4 \text{ mol}^2 \text{ m}^{-6} (1.0 \times 10^{-2} \text{ mol}^2 \text{ dm}^{-6}) \end{aligned} \quad (04 + 01)$$

(Note: Here  $K_c$  must be calculated from  $K_p$ , other answers are not accepted)

(iii) An extra amount of  $\text{CO}_2(\text{g})$  was added into the container described above at  $328^\circ\text{C}$ . When the equilibrium is re-established, the partial pressure of  $\text{CO}_2(\text{g})$  was four (4) times the partial pressure of  $\text{H}_2\text{O}(\text{g})$ . Calculate the partial pressures of  $\text{CO}_2(\text{g})$  and  $\text{H}_2\text{O}(\text{g})$  under this condition.

$$\text{Take, } P_{\text{H}_2\text{O}(\text{g})} = x \text{ Pa}$$

$$\therefore P_{\text{CO}_2(\text{g})} = 4x \text{ Pa}$$

$$\text{Now } K_p = P_{\text{H}_2\text{O}(\text{g})} \times P_{\text{CO}_2(\text{g})} = x \cdot 4x = 4x^2 \quad (05)$$

As the temperature is constant, (05)

$$2.5 \times 10^{11} \text{ Pa}^2 = 4x^2 \quad (05)$$

$$\left(\frac{2.5}{4}\right) \times 10^{11} \text{ Pa}^2 = x^2$$

$$\left(\frac{25}{4}\right) \times 10^{10} \text{ Pa}^2 = x^2$$


$$x = 2.5 \times 10^5 \text{ Pa}$$

$$\therefore P_{\text{H}_2\text{O}(\text{g})} = 2.5 \times 10^5 \text{ Pa} \text{ and} \quad (04 + 01)$$

$$P_{\text{CO}_2(\text{g})} = 1.0 \times 10^6 \text{ Pa} \quad (04 + 01)$$

**5(a): 75 marks**

(b) In order to determine the standard enthalpy change ( $\Delta H^\circ$ ) of the reaction,



the following experiment consisting of two steps (I and II) was carried out at room temperature.

**Step I:** 0.08 mol of  $\text{NaHCO}_3(\text{s})$  was added to  $100.00 \text{ cm}^3$  of  $1.0 \text{ mol dm}^{-3}$  HCl acid solution in a beaker. The maximum temperature fall was found to be  $5.0^\circ\text{C}$ .

[The reaction taking place:  $\text{NaHCO}_3(\text{s}) + \text{HCl}(\text{aq}) \rightarrow \text{Na}^+(\text{aq}) + \text{Cl}^-(\text{aq}) + \text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g})$ ]

**Step II:** 0.04 mol of  $\text{Na}_2\text{CO}_3(\text{s})$  was added to  $100.00 \text{ cm}^3$  of  $1.0 \text{ mol dm}^{-3}$  HCl acid solution in a beaker. The maximum temperature rise was found to be  $3.5^\circ\text{C}$ .

[The reaction taking place:  $\text{Na}_2\text{CO}_3(\text{s}) + 2\text{HCl}(\text{aq}) \rightarrow 2\text{Na}^+(\text{aq}) + 2\text{Cl}^-(\text{aq}) + \text{H}_2\text{O}(\text{l}) + \text{CO}_2(\text{g})$ ]

Specific heat capacity at constant pressure and density of HCl acid solution are  $4.0 \text{ J g}^{-1} \text{ K}^{-1}$  and  $1.0 \text{ g cm}^{-3}$  respectively. Assume that the changes in volume and density of the solutions after the addition of solids in the above two steps are negligible.

(i) Calculate the enthalpy changes (in  $\text{kJ mol}^{-1}$ ) of the reactions given in steps I and II above.

### Step I:

For the reaction with 0.08 moles of  $\text{NaHCO}_3(\text{s})$



$$Q = ms\theta \text{ Or } Q = mc\theta \quad (05)$$

$$= 100 \text{ g} \times 4.0 \text{ J g}^{-1} \text{ K}^{-1} \times 5 \text{ K} = 2000 \text{ J} = 2.0 \text{ kJ} \quad (04 + 01)$$

$\therefore$  for 1 mol

$$Q = 2.0 \text{ kJ} / 0.08 \text{ mol} \quad (05)$$

$$Q = \Delta H = + 25 \text{ kJ mol}^{-1} \text{ (heat loss)} \quad (04 + 01)$$

(02 marks for the sign)

**Step III:**

For the reaction with 0.04 moles of  $\text{NaHCO}_3(s)$



$$Q = ms\theta$$

$$= 100 \text{ g} \times 4.0 \text{ J g}^{-1} \text{ K}^{-1} \times 3.5 \text{ K} = 1400 \text{ J} = 1.4 \text{ kJ} \quad (04 + 01)$$

$\therefore$  for 1 mol

$$Q = 1.4 \text{ kJ} / 0.04 \text{ mol}$$


$$Q = \Delta H = -35 \text{ kJ mol}^{-1} \text{ (heat gain)} \quad (04 + 01)$$

(02 marks for the sign)

(II) Using the values obtained in (I) above and a thermochemical cycle, calculate  $\Delta H^\circ$  of the reaction



The reaction,  $2\text{NaHCO}_3(s) \rightarrow \text{Na}_2\text{CO}_3(s) + \text{CO}_2(g) + \text{H}_2\text{O(l)}$  can be written with the help of following thermochemical cycle,



(To award marks for the thermocycle balanced equations are necessary)

According to Hess's Law,  $\Delta H_1 = \Delta H_2 - \Delta H_3$  (05)

$$\Delta H_1 = 25 \times 2 - (-35) \text{ kJ mol}^{-1} \quad (05)$$

$$\Delta H_1 = +85 \text{ kJ mol}^{-1} \quad (04 + 01)$$

(III) State the condition under which a heat change of a reaction is equal to its enthalpy change.

At constant pressure. (05)

at 1 atm.

(IV) Identify two sources of error in the above experimental procedure.

Sources of errors

A calorimeter is not used/heat loss (heat loss to environment)

Dissolution enthalpies of solids are different

(03 x 2)

5(b): 75 marks

6. (a) (i) Explain why the rate of a reaction increases when the concentrations of the reactants are increased.

Increase in concentration causes an increase in the number of molecules in unit volume.

(05)

This increases the collision rate (Number of collisions in unit time). (05)

(ii) Give two reasons to explain why in general, the rate of a reaction increases with increasing temperature.

The collision rate and (05)


the fraction of reacting molecules having energy greater than the activation energy increase with increasing temperature. (05)

(iii) What is the relationship between order and molecularity of an elementary reaction?


Molecularity = Order of reaction. (10)

(iv) Sketch the structure of the activated complex of the elementary reaction,  $\text{NO} + \text{O}_2 \rightarrow \text{NO}_2 + \text{O}$ .

Label the bonds that are being formed as 'forming' and the bonds that are being broken as 'breaking'.



[Or]



(Correct structure 04 marks labelling 03 +03 )

(10)

(v) Write the rate expression for the elementary reaction,  $x\text{A} + y\text{B} \rightarrow z\text{C}$ , where the rate constant is  $k$ , and stoichiometric coefficients are  $x, y, z$ .

$$\text{Rate} = k[\text{A}]^x[\text{B}]^y \quad \underline{\text{or}} \quad \frac{1}{x} \left( \frac{-\Delta[\text{A}]}{\Delta t} \right) = k[\text{A}]^x[\text{B}]^y \quad \underline{\text{or}} \quad \frac{1}{x} \left( \frac{-d[\text{A}]}{dt} \right) = k[\text{A}]^x[\text{B}]^y$$

$$\underline{\text{or}} \quad \frac{-\Delta[\text{A}]}{\Delta t} = k[\text{A}]^x[\text{B}]^y \quad \underline{\text{or}} \quad \frac{-d[\text{A}]}{dt} = k[\text{A}]^x[\text{B}]^y \quad (10)$$

6(a): 50 marks

(b) The reaction  $xA + yB \rightarrow zC$  was studied in a two phase system consisting of an organic solvent and water. The compound A is soluble in both phases while compounds B and C are soluble only in the aqueous phase. The partition coefficient for the distribution of A between phases,  $K_D = \frac{[A]_{\text{org}}}{[A]_{\text{aq}}} = 4.0$ .

The compound A was added to the two phase system and allowed to reach equilibrium. The reaction was started by injecting the compound B to the aqueous phase. Temperature of the system was maintained at a constant value. The results of the experiments carried out are given below.

Experiment No.	Volume of the organic phase (cm <sup>3</sup> )	Volume of the aqueous phase (cm <sup>3</sup> )	Amount of A added to the system (mol)	Amount of B injected (mol)	Initial rate, $\left( \frac{-\Delta C_A}{\Delta t} \right)$ (mol dm <sup>-3</sup> s <sup>-1</sup> )
I	-	100.00	$1.00 \times 10^{-2}$	$1.00 \times 10^{-2}$	$1.20 \times 10^{-3}$
II	100.00	100.00	$1.25 \times 10^{-1}$	$1.00 \times 10^{-2}$	$7.50 \times 10^{-3}$
III	50.00	50.00	$6.25 \times 10^{-2}$	$1.00 \times 10^{-2}$	$1.50 \times 10^{-3}$

Note: Experiment I was carried out without the organic phase.

(i) Calculate the initial concentration of A in the aqueous phase in experiments I, II and III above.

#### Experiment (I)

$$\text{Concentration of A in the aqueous phase, } [A(I)]_{\text{aq}} = 1.00 \times 10^{-2} \text{ mol} / 100 \times 10^{-3} \text{ dm}^3 = 0.10 \text{ mol dm}^{-3}. \quad (01+01)$$

#### Experiment (II)

$$K = \frac{[A]_{\text{org}}}{[A]_{\text{aq}}} = 4$$

$$[A(II)]_{\text{aq}} = \frac{[A(II)]_{\text{org}}}{4.0} \quad (03)$$

Let x be the amount of A in the aqueous phase.

$$\frac{1.25 \times 10^{-1} \text{ mol} - x}{4 \times 100 \times 10^{-3} \text{ dm}^{-3}} = \frac{x}{100 \times 10^{-3} \text{ dm}^{-3}} \quad (02+01)$$

$$4x = 1.25 \times 10^{-1} \text{ mol} - x$$

$$x = 2.5 \times 10^{-2} \text{ mol} \quad (02+01)$$

$$[A(II)]_{\text{aq}} = \frac{x}{V} = \frac{2.5 \times 10^{-2} \text{ mol}}{100 \times 10^{-3} \text{ dm}^3}$$

$$[A(II)]_{\text{aq}} = 2.5 \times 10^{-1} \text{ mol dm}^{-3} \quad (02+01)$$

#### Experiment (III)

Let y be the amount of A in the aqueous phase.

$$[A(III)]_{\text{aq}} = \frac{y}{V}$$

$$\frac{y}{V} = \frac{6.25 \times 10^{-2} \text{ mol} - y}{V \times 4}$$

$$4y = 6.25 \times 10^{-2} \text{ mol} - y$$

$$y = \frac{6.25 \times 10^{-2} \text{ mol}}{5} = 0.0125 \text{ mol} \quad (02+01)$$

$$[A(III)]_{\text{aq}} = \frac{0.0125 \text{ mol}}{50.0 \times 10^{-3} \text{ dm}^3}$$

$$[A(III)]_{\text{aq}} = 2.5 \times 10^{-1} \text{ mol dm}^{-3} \quad (02+01)$$

(I) Find the order of the reaction with respect to A.

Finding the order with respect to A

In experiments (I) and (II), the concentration of B in the aqueous phase is the same.

$$1.2 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1} = k[0.1 \text{ mol dm}^{-3}]^x [0.1 \text{ mol dm}^{-3}]^y \quad (02 + 01)$$

$$7.5 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1} = k [0.25 \text{ mol dm}^{-3}]^x [0.1 \text{ mol dm}^{-3}]^y \quad (02 + 01)$$

$$1.5 \times 10^{-3} \text{ mol dm}^{-3} \text{ s}^{-1} = k [0.25 \text{ mol dm}^{-3}]^x [0.2 \text{ mol dm}^{-3}]^y \quad (02 + 01)$$

$$\frac{7.5 \times 10^{-5}}{1.2 \times 10^{-5}} = \left(\frac{0.25}{0.1}\right)^x \quad (03)$$

$$\frac{6.25}{1} = \left(\frac{2.5}{1}\right)^x$$

$$2.5^2 = 2.5^x$$

$$X=2 \quad (02)$$

(III) Find the order of the reaction with respect to B.

Finding the order with respect to B

In experiments II and III, the concentration of A in the aqueous phase is the same.

$$\frac{1.5 \times 10^{-3}}{7.5 \times 10^{-5}} = \left(\frac{0.2}{0.1}\right)^y \quad (03)$$

$$\frac{20}{1} = \left(\frac{2}{1}\right)^y$$

$$\log(20) = y \log 2$$

$$1.3010 = y \times .3010$$

$$Y = 4.32 = 4 \quad (02)$$

(IV) Calculate the rate constant of the reaction.

Finding the rate constant of the reaction.

Using data in experiment (I),

$$k = \frac{\text{Rate (I)}}{[A(I)]^2[B(I)]^4} \quad (03)$$

(4.32 can be used instead of 4)

$$k = \frac{1.20 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}}{(1.00 \times 10^{-3} \text{ mol dm}^{-3})^2 (1.00 \times 10^{-3} \text{ mol dm}^{-3})^4} \quad (02 + 01)$$

$$k = 1.2 \times 10^1 \text{ mol}^{-4} \text{ dm}^{12} \text{ s}^{-1} \quad (01)$$

(V) In the experiment III above, after adding A and allowing the system to reach equilibrium, if a volume of 10.00 cm<sup>3</sup> was removed from the organic phase, what can be stated about the initial rate of the reaction? Give reason/reasons for your answer.

The initial rate of the reaction will not change. The Concentration of A in the aqueous phase does not change.

(02 + 02)

6(b): 50 marks

- (c) A mixture of liquids X and Y behaves ideally. At a constant temperature, when the liquid phase is in equilibrium with the vapour phase in a closed rigid vessel contains 1.2 moles of X and 2.8 moles of Y, the total vapour pressure is  $3.4 \times 10^4$  Pa. At the same temperature, when the composition of the liquid phase is in equilibrium with the vapour phase is 1.2 moles of X and 4.8 moles of Y, the total vapour pressure is  $3.6 \times 10^4$  Pa. Calculate the saturated vapour pressures of X and Y at this temperature.

According to Raoult's law,  $p_i = x_i p_i^0$  (05)

$$P_{\text{total}} = P_x + P_y = X_x P_x^0 + X_y P_y^0 \quad (05)$$

For the first condition,

$$X_x = \frac{1.2 \text{ mol}}{1.2 \text{ mol} + 2.8 \text{ mol}} \quad (04 + 01)$$

$$X_x = 0.3 \text{ and therefore } X_y = 0.7 \quad (02 + 02 + 01)$$

For the second condition,

$$X_y = \frac{1.2 \text{ mol}}{1.2 \text{ mol} + 4.8 \text{ mol}} \quad (04 + 01)$$

$$X_y = 0.2 \text{ and therefore } X_x = 0.8 \quad (02 + 02 + 01)$$

Using the mole fractions, two equations can be constructed for the two conditions.

$$3.4 \times 10^4 \text{ Pa} = 0.3 P_x^0 + 0.7 P_y^0 \quad - (1) \quad (05)$$



$$3.6 \times 10^4 \text{ Pa} = 0.2 P_x^0 + 0.8 P_y^0 \quad - (2) \quad (05)$$

Solving the two equations,


$$P_x^0 = 2.0 \times 10^4 \text{ Pa} \quad (05) \quad (04 + 01)$$

$$P_y^0 = 4.0 \times 10^4 \text{ Pa} \quad (05) \quad (04 + 01)$$

6(c): 50 marks



(i) Show how you would prepare A and X in one more than five (5) steps in each case, starting X and Y as required.






(ii) Two compounds A and B are required to be prepared in the following:


\*  $\text{RCH}_2\text{Cl}$  is acceptable provided that the appropriate product is given.

\*  $(\text{CH}_3)_3\text{CCl}$  is unacceptable.

Notes: \* Instead of  $\text{CH}_3\text{Cl}$  any other appropriate allylic halide can be written.



7. (a) Show how you would carry out the following conversion in one more than five (5) steps.





30 marks

Note : The marks allocated for X and Y in the synthetic pathways can be awarded ONLY if the usage of X and Y is correct/appropriate.

7(b): 05 marks X 10 = 50 marks

(ii) Show how you would prepare the compound C using A and B given above, in not more than five (5) steps.




7(b)(ii): 05 marks X 8 = 40 marks


Note: Steps can be combined appropriately. Award marks appropriately for correct Answers.

No marks are awarded for Cu/300°C or heat

(c) Using your knowledge of the mechanism of the reaction between acetyl chloride and NaOH, propose a mechanism for the reaction between



and NaOH.



Notes: \*Steps can be combined appropriately. Award marks appropriately for correct Answers.

7(c): 30 marks

**(iii) Solution Y contains three cations**

**(Q)** The following tests were carried out to identify these cations.

	Test	Observation
①	Dilute HCl was added to a small portion of Y.	A white precipitate ( $P_1$ )
②	$P_1$ was separated by filtration and $H_2S$ was bubbled through the solution.	A black precipitate ( $P_2$ )
③	$P_2$ was separated by filtration. The filtrate was boiled to remove the $H_2S$ , cooled, and $NH_4OH/NH_4Cl$ was added.	No precipitate
④	$H_2S$ was bubbled through the solution.	A black precipitate ( $P_3$ )

**(Q)** The following tests were carried out for precipitates  $P_1$ ,  $P_2$  and  $P_3$ .

Precipitate	Test	Observation
$P_1$	I. Water was added to $P_1$ and the mixture was boiled. II. The mixture from I above was filtered while warm and the following tests were carried out on the filtrate ( $F_1$ ) and residue ( $E_1$ ).  Filtrate ( $F_1$ ) • Dilute $H_2SO_4$ was added to warm $F_1$ . Residue ( $E_1$ ) • $E_1$ was washed thoroughly with warm water and dilute $NH_4OH$ was added. • Thereafter, a KJ solution was added.	Part of $P_1$ dissolved.  A white precipitate $E_1$ dissolved. A dark yellow precipitate
$P_2$	$P_2$ was dissolved in warm dil. $HNO_3$ and a potassium chromate solution was added.	A yellow precipitate
$P_3$	I. $P_3$ was dissolved in warm conc. $HNO_3$ . II. The following were added to solution 1 above. • conc. HCl  • dil. $NH_4OH$	A pink coloured solution (solution 1)  A blue coloured solution (solution 2) A yellow-brown coloured solution (solution 3)

**(i) Identify the three cations. (Reasons are not required.)**



(08 x 3)

**(ii) Identify,**

I. precipitates  $P_1$ ,  $P_2$  and  $P_3$

II. species responsible for the colours of solutions, 1, 2 and 3.

(Note: Write chemical formulae only.)

I     $P_1$ :  $PbCl_2$  and  $AgCl$                    $P_2$ :  $PbS$                    $P_3$ :  $CoS$                   (05 x 4)

II    Solution 1:  $Co^{2+}$  OR  $Co(NO_3)_2$  OR  $[Co(H_2O)_6]^{2+}$                   (05)

Solution 2:  $[CoCl_4]^{2-}$                   (05)

Solution 3:  $[Co(NH_3)_6]^{2+}$                   (05)

(iii) Explain briefly why the cation/s that precipitate/s in ① ④ above does not/do not precipitate in acidic medium.

**For cations to precipitate from solution as sulfides**



In acidic medium,  $[S^{2-}]$  is reduced.



$K_{sp}$  of sulfides of Group IV cations >  $K_{sp}$  of sulfides of Group II cations (Mn<sup>2+</sup>, Zn<sup>2+</sup>, Ni<sup>2+</sup>, Co<sup>2+</sup>) (03)

As  $[S^{2-}]$  is reduced in acidic medium, it is not sufficient to precipitate the Group-IV cations as sulfides. Therefore, they remain in solution.

-103-

**8(a): 75 marks**

(b) A solid sample was found to contain  $(\text{NH}_4)_2\text{SO}_4$ ,  $\text{NH}_4\text{NO}_3$  and non-reactive substances. The following procedures were used to determine the amount of ammonium salts present in this sample.

A 1.00 g portion of the solid sample was dissolved in water and diluted to 250.00 cm<sup>3</sup> in a volumetric flask (hereafter referred to as solution S).

#### **Procedure 1**

A  $50.00 \text{ cm}^3$  portion of solution S was treated with an excess amount of a strong alkali ( $\text{NaOH}$ ) and the gas liberated was passed into  $30.00 \text{ cm}^3$  of  $0.10 \text{ mol dm}^{-3}$   $\text{HCl}$ . The volume of  $0.10 \text{ mol dm}^{-3}$   $\text{NaOH}$  required to neutralize the remaining  $\text{HCl}$  (using phenolphthalein as the indicator) was  $10.20 \text{ cm}^3$ .

### Procedure 2

To a  $25.00\text{ cm}^3$  portion of solution S, Al powder was added followed by an excess of strong alkali, and the mixture was heated. The gas liberated was passed into  $30.00\text{ cm}^3$  of  $0.10\text{ mol dm}^{-3}$  HCl. The volume of  $0.10\text{ mol dm}^{-3}$  NaOH required to neutralise the remaining HCl (using phenolphthalein as the indicator) was  $15.00\text{ cm}^3$ .


(Note: Completion of gas evolution in procedures 1 and 2 was checked using Ultrosil 2000.)

- (i) Identify the gas liberated in procedure 1.  
 (ii) Identify the gas liberated in procedure 2.  
 (iii) Write balanced chemical equations for the reactions taking place in procedures 1 and 2.  
 (iv) Calculate the mass percentage of each of the compounds  $(\text{NH}_4)_2\text{SO}_4$  and  $\text{NH}_4\text{NO}_3$ , in the solid sample. ( $\text{H} = 1$ ,  $\text{N} = 14$ ,  $\text{O} = 16$ ,  $\text{S} = 32$ )



(i)  $\text{NH}_3$  (03)

(ii)  $\text{NH}_3$  ~~hi~~ (13)

### (iii) Reactions in Procedure I



## Reactions in Procedure 2



**Note:** Since the question does not ask to write the reactions separately for procedures, students may write only the last three reactions as in the procedure 2. If so, award 07 marks each, to the reactions with NaOH (i.e. reactions 3 and 4).

(iv)	Amount of $(\text{NH}_4)_2\text{SO}_4$ in sample = $x$ g (01)	Amount of $\text{NH}_4\text{NO}_3$ in sample = $y$ g (01)
	Molar mass of $(\text{NH}_4)_2\text{SO}_4$ = 132 (01)	Molar mass of $\text{NH}_4\text{NO}_3$ = 80 (01)
	Moles of $(\text{NH}_4)_2\text{SO}_4$ = $\frac{x}{132}$ (01)	Moles of $\text{NH}_4\text{NO}_3$ = $\frac{y}{80}$ (01)

Procedure 1

Moles of NaOH =  $\frac{0.10}{1000} \times 10.20$  (01)

Therefore, moles of HCl reacted with NaOH =  $\frac{0.10}{1000} \times 10.20$  (01)

Therefore, moles of HCl reacted with  $\text{NH}_3$  =  $\frac{0.10}{1000} \times 30.0 - \frac{0.10}{1000} \times 10.20$  (01)

=  $\frac{0.10}{1000} \times 19.80$  (01)

Therefore, moles of  $\text{NH}_3(\text{g})$  liberated =  $\frac{0.10}{1000} \times 19.80$  (01)

Moles of  $(\text{NH}_4)_2\text{SO}_4$  in 50.00 cm<sup>3</sup> =  $\frac{x}{132} \times \frac{1}{5}$  (01)

Moles of  $\text{NH}_4\text{NO}_3$  in 50.00 cm<sup>3</sup> =  $\frac{y}{80} \times \frac{1}{5}$  (01)

Therefore, Moles of  $\text{NH}_3$  given out by 50.00 cm<sup>3</sup> of solution (Eq. (1) & (2)) =

$$\frac{x}{132} \times \frac{1}{5} \times 2 + \frac{y}{80} \times \frac{1}{5} = \frac{0.10}{1000} \times 19.80 \quad \dots \dots \dots \text{(6)} \quad \text{(07)}$$

Similarly for Procedure 2

Moles of NaOH =  $\frac{0.10}{1000} \times 15.00$  (01)

Therefore, moles of HCl reacted with NaOH =  $\frac{0.10}{1000} \times 15.00$  (01)

Therefore, moles of HCl reacted with  $\text{NH}_3$  =  $\frac{0.10}{1000} \times 30.0 - \frac{0.10}{1000} \times 15.00$  (01)

=  $\frac{0.10}{1000} \times 15.00$  (01)

Therefore, moles of  $\text{NH}_3(\text{g})$  liberated =  $\frac{0.10}{1000} \times 15.00$  (01)

$$\text{Moles of } (\text{NH}_4)_2\text{SO}_4 \text{ in } 25.00 \text{ cm}^3 = \frac{1}{132} \times \frac{1}{10} \quad (01)$$

$$\text{Moles of } \text{NH}_4\text{NO}_3 \text{ in } 25.00 \text{ cm}^3 = \frac{1}{80} \times \frac{1}{10} \quad (01)$$

Moles of NH<sub>3</sub> given out by 25.00 cm<sup>3</sup> of solution (Eq. (4), (6) & (9)) -

$$\frac{x}{132} \times \frac{1}{10} \times 2 + \frac{y}{80} \times \frac{1}{10} + \frac{y}{80} \times \frac{1}{10} = \frac{0.10}{1000} \times 15.00 \quad (7) \quad (01)$$

$$\frac{x}{132} \times \frac{1}{10} \times 2 + \frac{2y}{80} \times \frac{1}{10} = \frac{0.10}{1000} \times 15.00 \quad (8) \quad (01)$$

$$\frac{x}{132} \times \frac{1}{5} + \frac{y}{80} \times \frac{1}{5} = \frac{0.10}{1000} \times 15.00 \quad (9) \quad (01)$$

$$\text{Eq. (6)} \quad \frac{x}{132} \times \frac{1}{5} + \frac{y}{80} \times \frac{1}{10} = \frac{0.10}{1000} \times 19.80 \times \frac{1}{2} \quad (6)$$

$$\text{Eq. (8)} - \text{Eq. (6)} \quad \frac{y}{80} \times \frac{1}{10} = \frac{0.10}{1000} \times 15.00 - \frac{0.10}{1000} \times 9.90 \quad (01)$$

$$y = \frac{0.10}{1000} \times 5.10 \times 800; \quad (02)$$

$$y = 0.408 \quad (02)$$

Substituting  $y = 0.408$  in Eq.(8) ( $y=0.408$  can also be substituted to Eq. (6))

$$\begin{aligned} \frac{x}{132} \times \frac{1}{5} + \frac{0.408}{80} \times \frac{1}{5} &= \frac{0.10}{1000} \times 15.00 \\ \frac{x}{660} &= \frac{0.10}{1000} \times 15.00 - \frac{0.408}{400} = \frac{(6.0 - 4.08)}{4000} \\ x &= 0.317 \end{aligned} \quad (02)$$

$$\% \text{ of } (\text{NH}_4)_2\text{SO}_4 = \frac{0.317}{1.0} \times 100 = 31.7\% \quad (05)$$

$$\% \text{ of } \text{NH}_4\text{NO}_3 = \frac{0.408}{1.0} \times 100 = 40.8\% \quad (05)$$

8(b)(iv): 75 marks

**Alternative Answer 1**

### Procedure 1

Amount of  $\text{NH}_3$  collected =  $(30.00 \times 0.10 - 10.20 \times 0.10) \times 10^{-3} \text{ mol}$  (02)

$$= 1.98 \times 10^{-3} \text{ mol}$$

$$\text{Amount of NH}_3 \text{ from 1 g of sample} = 1.98 \times 10^{-3} \times 5$$

$$= 9.90 \times 10^{-3} \text{ mol}$$

#### **Procedure 2**

$$\text{Amount of NH}_3 \text{ collected} = (30.00 \times 0.10 - 15.00 \times 0.10) \times 10^{-3} \text{ mol} \quad (02)$$

$$= 1.50 \times 10^{-3} \text{ mol}$$

$$\text{Amount of NH}_3 \text{ from 1 g of sample} = 1.50 \times 10^{-3} \times 10 \text{ mol}$$

$$\approx 15.0 \times 10^{-3} \text{ mol}$$

**Number of moles of  $\text{NH}_4\text{NO}_3$  in 1 g of sample = x**

Number of moles of  $(\text{NH}_4)_2\text{SO}_4$  in 1 g of sample

In Procedure 1, NH<sub>3</sub> from NH<sub>4</sub>NO<sub>2</sub> and (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> is captured.

$$\text{Therefore, } x + 2y = 9.90 \times 10^{-3} \text{ mol} \quad (1) \quad (67)$$

In Procedure 2,  $\text{NO}_3^-$  is converted to  $\text{NH}_4^+$  and evolved together with  $\text{NH}_3$  from  $\text{NH}_4\text{NO}_3$  and  $\text{NH}_4\text{Cl}$ .

$$\text{Therefore, } 2x + 2y = 15.0 \times 10^{-3} \text{ mol} \quad (2) \quad (cont)$$

$$(2) - (1) \quad x = (15.0 - 2.0) \times 10^3 = 13$$

$$\approx 5.1 \times 10^{-3} \text{ mol}$$

From (1)  $y = \frac{1}{2}(9.9 - 5.1) \times 10^{-3}$  (02)

$$= 2.4 \times 10^{-3} \text{ mol} \quad (02)$$

$$\% \text{ of } \text{NH}_4\text{NO}_3 \text{ in sample} = \frac{5.1 \times 10^{-3} \text{ mol} \times 80 \text{ g/mol} \times 100}{1.0 \text{ g}} \quad (\text{04 + 01 for mol. mass})$$

= 40.8 % (05)

$$\begin{aligned} \text{\% of } (\text{NH}_4)_2\text{SO}_4 \text{ in sample} &= \frac{2.4 \times 10^{-3} \text{ mol} \times 132 \text{ g mol}^{-1} \times 100}{1.0 \text{ g}} \quad (04 + 01) \\ &\quad (04 + 01 \text{ for mol. mass}) \\ &= 31.7\% \quad (05) \end{aligned}$$

**Alternative Answer 2****Procedure 1**


Initial number of moles of HCl =  $\frac{0.10}{1000} \times 30.00$  (01)

Number of moles of HCl remaining =  $\frac{0.10}{1000} \times 10.20$  (01)

HCl : NaOH = 1: 1

Therefore, number of moles of HCl used =  $\frac{0.10}{1000} \times 30.00 - \frac{0.10}{1000} \times 10.20$  (02)

$$= \frac{0.10}{1000} \times 19.80 = 0.00198 \quad (02)$$



Therefore, number of moles of NH<sub>3</sub> evolved (from NH<sub>4</sub>NO<sub>3</sub> & (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, in 50.0 cm<sup>3</sup> of solution) = 0.00198 (01)

**Procedure 2**

Initial number of moles of HCl =  $\frac{0.10}{1000} \times 30.00$  (01)

Number of moles of HCl remaining =  $\frac{0.10}{1000} \times 15.00$  (01)

Therefore, number of moles of HCl used =  $\frac{0.10}{1000} \times 30.0 - \frac{0.10}{1000} \times 15.00$  (02)

$$= \frac{0.10}{1000} \times 15.00 \quad (01)$$

$$= 0.0015$$

Number of moles in 50.00 cm<sup>3</sup> =  $0.0015 \times 2 = 0.003$  (01)

Therefore, number of moles of NO<sub>3</sub><sup>-</sup> in 50.00 cm<sup>3</sup> = 0.003 - 0.00198 = 0.00102 (07)

Mass of NH<sub>4</sub>NO<sub>3</sub> in 250.0 cm<sup>3</sup> =  $0.00102 \times 80 \times 5 \text{ g}$  (01)

% of NH<sub>4</sub>NO<sub>3</sub> in sample =  $\frac{0.00102 \text{ mol} \times 80 \text{ g mol}^{-1} \times 100}{1.0 \text{ g}}$  (04 + 01 for mol. mass)  
= 40.8% (05)

Number of moles of NH<sub>3</sub> obtained from NH<sub>4</sub>NO<sub>3</sub> (from 50.00 cm<sup>3</sup>) = 0.00102

Therefore, number of moles of NH<sub>3</sub> obtained from (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> = 0.00198 - 0.00102 = 0.00096 (07)

(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> : NH<sub>3</sub> = 1:2

Therefore, number of moles of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> =  $\frac{0.00096}{2} = 0.00048$  (01)

Mass of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> in sample =  $0.00048 \times 132 \times 5 \text{ g}$  (01)

% of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> in sample =  $\frac{0.00048 \text{ mol} \times 132 \text{ g mol}^{-1} \times 5 \times 100}{1.0 \text{ g}}$  (04 + 01 for mol. mass)  
= 31.7% (05)

8(b): 75 marks

9. (a) Consider the following industrial processes.

- I. Manufacture of bleaching powder
- II. Manufacture of calcium carbide
- III. Manufacture of urea
- IV. Manufacture of sulphuric acid (contact process)

(i) State the starting materials used in each process.




- I Bleaching powder –  $\text{CaCO}_3$  (lime stone)/ $\text{CaO}$  (quick lime),  $\text{Ca}(\text{OH})_2$  (slaked lime) and  $\text{Cl}_2$  gas
- II Calcium carbide –  $\text{CaO}$  (quick lime),  $\text{CaCO}_3$  (lime stone), and coke (C)
- III Urea –  $\text{NH}_3$  (l or Gas) and  $\text{CO}_2$  (l or Gas)
- IV  $\text{H}_2\text{SO}_4$  – sulfur (s)/ $\text{FeS}_2$  (iron pyrite), air and  $\text{H}_2\text{O}$

(02 X 9 = 18 marks)

9(a)(i): 18 marks

(ii) Write balanced chemical equations for the reactions taking place in each process, along with proper conditions wherever necessary.

I Bleaching powder



9(a)(ii)I: 11 marks

II Calcium carbide




OR





OR



9(a)(ii)II: 05 marks



**IV      Sulphuric acid**

[9(a)(ii)IV: 15 marks]

[9(a)(ii): 41 marks]

Note : Physical states not required for a(I) and (II)

- (ii) State two uses for each of the following:  
bleaching powder, calcium carbide, urea and sulphuric acid

UsesBleaching powder

- Bleaching agent
- Disinfects water

Calcium carbide

- To induce flowering
- To ripen fruits
- Production of oxyacetylene flame
- Produces acetylene

Urea

- To manufacture nitrogen rich fertilizers
- Manufacture of urea-formaldehyde polymer

 $\text{H}_2\text{SO}_4$ 

- Manufacture of phosphate fertilizer
- Manufacture of  $(\text{NH}_4)_2\text{SO}_4$
- Manufacture of rayon and plastics
- Manufacture of dyes
- Manufacture of explosives
- Manufacture of drugs
- Battery acid
- To dry gases (e.g.  $\text{Cl}_2$  gas)
- 

$(02 + 02) \times 4 = 16 \text{ marks}$

Note: For Uses - Award marks for any acceptable uses not listed in the marking scheme after getting approval from Chief Examiner.

9(a)(iii): 16 marks

9(a): 75 marks

(b) The major environmental problems we are facing at present are ozone layer depletion (OLD), global warming (GW) and acid rain (AR). The questions given below are related to the environment and the problems mentioned above.

(i) The carbon and nitrogen cycles are two important chemical cycles that operate in the environment.

I. With regard to the carbon cycle, state one main form of carbon that is present in each of the following:

atmosphere, plants, water, Earth's crust

II. State briefly how  $N_2$  gas in the atmosphere is removed and replenished in the nitrogen cycle.

III. State two ways by which microorganisms participate in the carbon cycle.

I

- Atmosphere –  $CO_2$  (02)
- Plants – glucose/ starch/ cellulose/ lignin (02)
- Water –  $HCO_3^- / CO_3^{2-} (CO_2(aq))$  (02)
- Earth's crust –  $CaCO_3 / CaCO_3.MgCO_3 /$  fossil fuel (coal, petroleum) / graphite (02)

II Removed via

- Industrial fixation (Haber process)/internal combustion engine/high temperature combustion (02)
- Biological fixation (nitrogen fixing bacteria) (02)
- Atmospheric fixation (by lightning) (02)


Replenished via

- Denitrifying bacteria that converts  $NO_3^-$  to  $N_2$  gas (02)

III Under aerobic conditions carbon compounds in soil are converted to  $CO_2$  (02)  
whereas, under anaerobic conditions they are converted to  $CH_4 & CO$  (02)

9(b)(i): 20 marks

(ii) Identify the two main nitrogen containing compounds present in the atmosphere that contribute to acid rain. With the aid of balanced chemical equations show how these compounds make rain water acidic.



$HNO_3$  is a strong acid which gives  $H^+ + NO_3^-$  OR  $(HNO_3 \rightarrow H^+ + NO_3^-)$  (02)

9(b)(ii): 10 marks

(iii) Identify two contributing industrial processes per environmental problem stated above (OLD, GW, AR). Identify one chemical compound that is liberated to the atmosphere by each of these industrial processes.

### Ozone layer depletion

- Air-conditioning industry (CFC / HCFC / aerosol)
  - Refrigerator Industry (CFC / HCFC)
  - Aircraft Industry or supersonic jets (NO<sub>2</sub>)
  - Agrochemical Industry (CH<sub>3</sub>Br Weedicide, fumigant)
  - Fire extinguisher industry using or releasing Cl<sub>2</sub>

96

**(Industry (02) Chemical species (01)) X 2**

### **Global warming**

- Agriculture ( $\text{CH}_4$ ,  $\text{N}_2\text{O}$ )
  - Air-conditioners (CFC/ HCFC)
  - Refrigerator industry (CFC / HCFC)
  - Poultry farming ( $\text{CH}_4$ )
  - Coal power plants ( $\text{CO}_2$ )
  - Nitrate fertilizers
  - Agrochemical industry ( $\text{CH}_3\text{Br}$  Weedicide, fumigant)
  - Halons in fire extinguishers
  - Any industry producing / using chlorine gas
  - Aerosol / spraying agents

8k-2

**(Industry (02) Chemical species (01)) X 2**

### **Acid rain**

- Burning fossil fuels in transportation ( $\text{NO}_x, \text{NO}_2$ )
  - Burning of fossil fuel in coal power plants ( $\text{SO}_2$ )
  - High temperature combustion ( $\text{NO}_x, \text{NO}_2$ )

96-

(Industry (02) Chemical species (01) ) X 2

9(b)(ii): 18 marks


(iv) Identify the main industrial process that contributes significantly to the addition of nitrogen compounds to water and soil. Comment on the pathways by which these compounds are added to water and soil.

### ~~Haber Process (NH<sub>3</sub> Production)~~

(cont)

~~NH<sub>3</sub> can be converted to fertilizers~~


103



**HNO<sub>3</sub>** is used to make fertilizers (O<sub>2</sub>) such as KNO<sub>3</sub> (O<sub>2</sub>), NH<sub>4</sub>NO<sub>3</sub> (O<sub>2</sub>), urea (O<sub>2</sub>), which are added to soil.

9

II Liquid NH<sub>3</sub> is directly used as a fertilizer. (01)



**9(b)(iv): 18 marks**

(v) Inappropriate disposal of solid municipal waste as in the Meethotamulla event contributes significantly to one of the three environmental issues stated above. Identify this environmental issue and state briefly how the disposal of solid municipal waste contributes to it.

As a result of the unregulated disposal of solid waste, large amounts of methane (02) are produced (between the layers of solid waste) due to the action of bacteria (01) under anaerobic conditions (02). This methane is a greenhouse gas (01) and contributes to global warming (03). 9(b)(v): 09 marks

9 (b) (v): 09 marks

(iv) Using the  $E^\circ$  values given, calculate the electromotive force of the cell.

Cell e.m.f

$$E^{\text{cell}} = E^\circ_{\text{Hg/Hg}_2\text{Cl}_2} - E^\circ_{\text{Ag/AgCl}}$$

$$= 0.27\text{V} - 0.22\text{V}$$

(04+01)

$$= 0.05\text{V}$$

(04+01)

(v) Give the standard cell notation of the above electrochemical cell.

Cell notation



[OR]

(10)



(Note: Do not deduct marks if Pt is written)

Note : For parts (i) to (v), Physical states are required

(vi) Does the electromotive force of the above electrochemical cell depend on the chloride ion concentration? Give reason/s for your answer.

The cell potential does not depend on the chloride ion concentration. Chloride ion concentration is not present in the cell reaction. (05 + 05)

(vii) Calculate the change in the mass of Ag(s) + AgCl(s), when a current of 0.10A was drawn for a period of 60 minutes from the cell.

A current of 0.10A was drawn for 60 min.

The amount of charged passed through the cell =  $0.10\text{A} \times 60 \text{ min} \times 60 \text{ s min}^{-1}$

$$= 360 \text{ C}$$

(04 + 01)

Electrode reaction in the left electrode Ag(s) + Cl<sup>-</sup> (aq)  $\rightarrow$  AgCl(s).

For every electron passed through the cell, one Ag atom combines with Cl<sup>-</sup> to form AgCl(s).

Weight gain of Ag + AgCl = mass of Cl<sup>-</sup> combined with Ag.

$$= 35.5 \text{ g mol}^{-1} \times 360 \text{ C} / 96500 \text{ C mol}^{-1}$$

(04 + 01)

$$= 0.132 \text{ g}$$

(04 + 01)

(viii) What would be the chloride ion concentration in the solution after drawing the current in (vii) above?

Chloride ion concentration does not change or chloride ion concentration = 1.0 mol dm<sup>-3</sup>.

(10)

10(b): 75 marks